

Proof of a conjectured q, t -Schröder identity

William J. Keith

Drexel University, 3141 Chestnut, Philadelphia, PA 19104, USA
wj26@drexel.edu

January 22, 2010

Abstract

A conjecture of Chunwei Song on a limiting case of the q, t -Schröder theorem is proved combinatorially. The proof matches pairs of tableaux to Catalan words in a manner that preserves differences in the maj statistic.

1 Introduction

In [1], Chunwei Song examined the combinatorics of a limiting case of the q, t -Schröder Theorem, proved by Haglund in [2]. To state the main conjecture which concerns this paper, use the standard notation

$$(a; q)_n = (1 - a)(1 - aq) \dots (1 - aq^{n-1}), \quad \begin{bmatrix} n \\ k \end{bmatrix} = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}$$

and let $S_{n,i}$ be the set of permutations on n letters with longest increasing subsequence of length i . Define

$$F_{n,k} := \sum_{i=1}^k \sum_{\substack{a_1 + \dots + a_k = n \\ a_i > 0}} q^{\sum_{i=1}^k \binom{a_i}{2}} t^{\sum_{i=1}^{k-1} (k-i)a_i} \frac{1}{(t^k; q)_{a_1} (q, q)_{a_k}} \\ \times \prod_{i=1}^{k-1} \begin{bmatrix} a_i + a_{i+1} - 1 \\ a_i \end{bmatrix} \frac{1}{(t^{k-i}; q)_{a_i + a_{i+1}}} \times (q; q)_n (t; t)_n$$

and, with maj the major index of a permutation,

$$H_{n,k}(q, t) := \sum_{i=1}^k \sum_{\sigma \in S_{n,i}} q^{\text{maj}(\sigma)} t^{\binom{n}{2} - \text{maj}(\sigma^{-1})}.$$

Then Song conjectured, for the $q = t$ case,

Conjecture 1 $F_{n,k}(q, q) = H_{n,k}(q, q)$.

For $k = 1$, both sides are $q^{\binom{n}{2}}$. For $k > 1$, the conjecture remains. Song rewrote the $k = 2$ case as (Conjecture 3.7 in [1]):

Conjecture 2

$$\sum_{w \in CW_n} q^{\text{maj}(w) - n \text{des}(w)} = \sum_{\sigma \in S_n(123)} q^{\text{maj}(\sigma) - \text{maj}(\sigma^{-1})}.$$

Here $S_n(123)$ is the set of 123-avoiding permutations (those with longest increasing subsequence of length no more than 2), and CW_n is the set of *Catalan words*: words $x_1 \dots x_{2n}$ with $x_i \in \{0, 1\}$ such that there are exactly n 1s and no initial segment has more 1s than 0s. The *descent set* of such a word is $\text{Des}(w) = \{i | x_i = 1, x_{i+1} = 0\}$. The *maj* of such a word is $\sum_{x_i \in \text{Des}(w)} x_i$ and $\text{des}(w) = |\text{Des}(w)|$. Song proved a subset of this conjecture, proving equality for the set of w and σ representing involutions, where w is an involution if $w = w^{-1}$ under the inversion operation that reverses w and exchanges 0 and 1, an operation that corresponds to reversing the associated Dyck path. This article proves the full $k = 2$ case by a bijection that may be generalizable to any k .

2 The bijection

Since $\text{maj}(w^{-1}) = 2n \text{des}(w) - \text{maj}(w)$ (from the proof of Theorem 3.10 in [1]), we can rewrite Conjecture 2 more symmetrically as

Theorem 1

$$\sum_{w \in CW_n} q^{\text{maj}(w) - \text{maj}(w^{-1})} = \sum_{\sigma \in S_n(123)} q^{2(\text{maj}(\sigma) - \text{maj}(\sigma^{-1}))}.$$

A map that takes Catalan words of length $2n$ to permutations $\sigma \in S_n(123)$ by a mapping $\phi(w) = \sigma$ such that $\text{maj}(w) - \text{maj}(w^{-1}) = 2(\text{maj}(\sigma) - \text{maj}(\sigma^{-1}))$ would now prove the theorem. This we construct.

Define a *Catalan half-word* of length n to be a 01 word $x_1 \dots x_n$ with no initial segment having more 1s than 0s. The number of 1s may vary, though the initial-segment condition means there are never more than $\lfloor \frac{n}{2} \rfloor$. The set of ordered pairs (w_1, w_2) of Catalan half-words of length n in which w_1 and w_2 have the same number of 1s is in bijection with CW_n by $(w_1, w_2) \rightarrow w_1 w_2^{-1}$, inversion being the same as before (reversal and exchanging 0 and 1) and juxtaposition denoting simple concatenation. Clearly (w_2, w_1) corresponds to the inverse of (w_1, w_2) .

A *tableaux*, or standard Young tableaux, of n boxes is an arrangement of the numbers 1 through n on points of the lattice in the first quadrant such that rows increase when read left to right and columns increase when read from the bottom up, and there are no gaps between occupied lattice points in rows or columns. The set of occupied lattice points is the shape of such a tableaux. The descent

set $Des(\tau)$ of a tableaux τ is the set of i with $i + 1$ in a higher row than i in τ , and $\text{maj}(\tau)$ is the sum of such i . Pairs of tableaux (τ_1, τ_2) of n boxes wherein both have the same shape are in bijection with permutations $\sigma \in S_n$ through the famous Robinson-Schensted correspondence, which has the properties that pairs (τ_2, τ_1) correspond to σ^{-1} (so that diagonal pairs (τ_1, τ_1) correspond to involutions), tableaux with exactly i columns correspond to permutations in $S_{n,i}$, and $\text{maj}(\tau_2) = \text{maj}(\sigma)$.

Lemma 1 *The set T_D of tableaux τ of n boxes, with no more than 2 columns and $j \geq 0$ entries in the second column, that have $D = Des(\tau) = \{i_1, \dots, i_k\}$, is equal in cardinality to the set CW_D of Catalan half-words w of length n that have j 1s and patterns $x_i x_{i+1} = 01$ where $n - i \notin D$.*

Example 1 *The two tableaux with descent set $D = \{1, 3, 4, 6\}$, $n = 7$, $j = 3$ are*

$$\begin{array}{|c|c|} \hline 7 & \\ \hline 5 & 6 \\ \hline 2 & 4 \\ \hline 1 & 3 \\ \hline \end{array} = \tau_1, \quad \begin{array}{|c|c|} \hline 5 & \\ \hline 4 & 7 \\ \hline 2 & 6 \\ \hline 1 & 3 \\ \hline \end{array} = \tau_2.$$

The permutation σ associated to the pair (τ_1, τ_2) by the Robinson-Schensted correspondence is (7562143), with maj 14, and its inverse is (5476231), also with maj 14, so their difference is 0. The two words in $CW_{\{2,5\}}$ for $n = 7$, $j = 3$ are $w_1 = 0010011$ and $w_2 = 0011010$. The Catalan word w associated to the ordered pair (w_1, w_2) is $w = 00100111010011$. Then $w^{-1} = 00110100011011$, and $\text{maj}(w) - \text{maj}(w^{-1}) = 3 + 8 + 10 - 4 - 6 - 11 = 0$.

Proof of Lemma 1

Select n , j , and $D = \{i_1, \dots, i_k\}$. These choices fix the following parts of the tableaux and Catalan half-words under consideration:

$$\overbrace{0 \cdots 00 \quad \underbrace{01}_{n-i_k, n-i_k+1} \quad 11 \cdots 100 \cdots 0 \quad \underbrace{01}_{n-i_{k-1}, n-i_{k-1}+1} \quad \cdots \quad \underbrace{01}_{n-i_1, n-i_1+1} \quad 11 \cdots 100 \cdots 0}^n$$

i_k	
$\vdots$	
i_2	$\vdots$
$\vdots$	$i_k + 1$
i_1	$\vdots$
$\vdots$	$i_2 + 1$
2	$\vdots$
1	$i_1 + 1$

In the word, the segments between 01 entries must be filled with nonincreasing sequences: $00 \cdots 0$, $11 \cdots 10 \cdots 0$, or $11 \cdots 1$. The choice of how to fill each segment is thus completely determined by the number of 1s we choose to enter. This choice is bounded by the initial-segment condition, the number of spaces in each segment, and the total number of 1s available ($j - k$).

In the tableaux, the sections between $i_a + 1$ and i_{a+1} must be filled without producing any more descents. The only way to do this is by listing the entries from $i_a + 2$ in the right-hand column up to some $i_a + c$, and then listing the remaining entries from $i_a + c + 1$ to $i_{a+1} - 1$ in the left-hand column. The choice of how to fill each section is thus completely determined by the choice of c . This choice is bounded by the condition that the two parts of the fill cannot overlap, the length of the interval between i_a and i_{a+1} , and the total number of spaces available in the right-hand column ($j - k$).

The two sets of conditions are exactly analogous. We can fill the tableaux section between $i_1 + 1$ and i_2 by choosing a segment of length up to $i_1 - 1$ entries in the right-hand column, unless bounded by the spaces available. We can fill the word segment between the last descent and the end of the word with a number of 1s up to the space available, $i_1 - 1$ spaces, unless bounded by the number of 1s available. Fill each segment up the tableaux and back along the word, making equal legal choices at each step. $\square$

This correspondence defines our map. Let $\phi : S_n(123) \rightarrow CW_n$ by sending $\sigma \rightarrow (\tau_1, \tau_2) \rightarrow (w_1, w_2)$, where the first map is the Robinson-Schensted correspondence and the second sends the tableaux τ_i to the word w_i such that w^{-1} has the same descent set as τ_i and the number of 1s in the segments between 01 entries at positions $n - i_{a-1}$ and $n - i_a$ are given by the number of elements in the right-hand column of τ_1 above each $i_a + 1$.

Example 2 The tableaux τ_1, τ_2 and words w_1, w_2 from Example 1 are from corresponding T_D and CW_D . The fixed and selectable sections in the words of CW_D look like $[0][01][?][01][?]$. Since in τ_1 the entry 4 appears above 3 in the

right-hand column, the map assigns $\phi(\tau_1) = 0010011 = w_1$, filling the segment after position $n - i_1$ with a 1. Since in τ_2 the entry 4 appears in the left-hand column, and the 7 appears in the right-hand column after $i_2 + 1 = 6$, the map fills the segment right of $n - i_2 = n - 5$ with the 1, giving $\phi(\tau_2) = 0011010 = w_2$.

All tableaux with the same descent set have the same maj. (Other tableaux may have the same maj statistic; there are as many such sets as there are partitions of $\text{maj}(w)$ into distinct nonconsecutive parts less than n .) As can be seen above, not all of the words in CW_D have the same maj; instead, it will be shown in the next section that each makes the same contribution to $\text{maj}(w) - \text{maj}(w^{-1})$ under the correspondence of Catalan half-words with Catalan words described above. To complete the proof of the theorem it remains only to show that $\text{maj}(\tau_i)$ and $\text{maj}(w_i)$ are related as claimed.

3 Proofs of required properties

Lemma 2 *Let w_1 and w_2 be two words in some CW_D , thus having the same length n , the same number of 1s j , and the same locations of 01 patterns starting at $\{n - i_1, \dots, n - i_k\}$. Let w_3 be any other Catalan half-word with j 1s. Say $w_b = w_1 w_3^{-1}$ and $w_c = w_2 w_3^{-1}$. Then $\text{maj}(w_b) - \text{maj}(w_b^{-1}) = \text{maj}(w_c) - \text{maj}(w_c^{-1})$.*

Proof If w_1 and w_2 are different, there will be an earliest digit in which they differ, say x_i , with w_1 having an entry of 0 and w_2 having an entry of 1. Since by hypothesis they possess the same number of 1s, there will be a latest digit, say x_j , in which w_1 has an entry of 1 and w_2 has an entry of 0. Suppose we alter w_1 by exchanging its 0 and 1 entry in these two places. Call w_4 the resulting word. (The initial-segment condition may be violated in w_4 , but the maj statistic is defined for all 01-words.) What is the relation between $\text{maj}(w_1 w_3^{-1}) - \text{maj}((w_1 w_3^{-1})^{-1})$ and $\text{maj}(w_4 w_3^{-1}) - \text{maj}((w_4 w_3^{-1})^{-1})$?

$$\begin{aligned} w_b &= w_1 w_3^{-1} = 0 \cdots x_{i-1} 0 x_{i+1} \cdots x_{j-1} 1 x_{j+1} \cdots w_3^{-1} \\ w_b^{-1} &= w_3 w_1^{-1} = w_3 \cdots (1 - x_{j+1}) 0 (1 - x_{j-1}) \cdots (1 - x_{i+1}) 1 (1 - x_{i-1}) \cdots 1 \\ w_4 w_3^{-1} &= 0 \cdots x_{i-1} 1 x_{i+1} \cdots x_{j-1} 0 x_{j+1} \cdots w_3^{-1} \\ w_4 w_3^{-1} &= w_3 \cdots (1 - x_{j+1}) 1 (1 - x_{j-1}) \cdots (1 - x_{i+1}) 0 (1 - x_{i-1}) \cdots 1 \end{aligned}$$

Note that x_{j+1} may be the first digit of w_3^{-1} . Additionally, it does not matter to $\text{maj}(w) - \text{maj}(w^{-1})$ whether the transition between w_1 and w_3^{-1} is a descent, or not; if it is, then w_1 ends in a 1 and the first digit of w_3^{-1} is a 0, and so the last digit of w_3 is 1 and the first digit of w_1 is 0 as well. Likewise, if the transition is not a descent in w , it is not in w^{-1} either.

Because w_1 and w_2 are both in the same CW_D , we cannot have 01 patterns in different places in the two words. This proscribes the case where $x_{i-1} = 0$, since then $x_{i-1} x_i$ would be 00 in w_1 , but would be a 01 sequence in w_2 . Similarly, we cannot have $x_{i+1} = 1$ (in particular, we cannot have $x_j = x_{i+1}$), $x_{j-1} = 0$,

or $x_{j+1} = 1$ with the exception of the possibility that x_{j+1} is the first digit of w_3^{-1} . If it is not, then our situation is

$$\begin{aligned} w_b &= w_1 w_3^{-1} = 0 \cdots 100 \cdots 110 \cdots w_3^{-1} \\ w_b^{-1} &= w_3 w_1^{-1} = w_3 \cdots 100 \cdots 110 \cdots 1 \\ w_4 w_3^{-1} &= 0 \cdots 110 \cdots 100 \cdots w_3^{-1} \\ w_3 w_4^{-1} &= w_3 \cdots 110 \cdots 100 \cdots 1 \end{aligned}$$

in the case when x_{j+1} is not the first digit of w_3^{-1} . We have descents in w_b at positions x_{i-1} and x_j , and in w_b^{-1} at positions x_{2n-j} and x_{2n-i+1} , contributing to $\text{maj}(w_b) - \text{maj}(w_b^{-1})$ a total of $i-1+j-(2n-j+2n-i+1) = 2i+2j-4n-2$. In $w_4 w_3^{-1}$, the first descent is moved forward a place, and the second is moved back 1, and likewise for the descents in the inverse. Thus the total does not change.

If x_{j+1} is the first digit of w_3^{-1} , then our situation is

$$\begin{aligned} w_b &= w_1 w_3^{-1} = 0 \cdots 100 \cdots 11 w_3^{-1} \\ w_b^{-1} &= w_3 w_1^{-1} = w_3 00 \cdots 110 \cdots 1 \\ w_4 w_3^{-1} &= 0 \cdots 110 \cdots 10 w_3^{-1} \\ w_3 w_4^{-1} &= w_3 10 \cdots 100 \cdots 1 \end{aligned}$$

If $x_{j+1} = 1$, then the last digit of w_3 is 0. Then w_b has a descent at x_{i-1} and w_b^{-1} a descent at x_{2n-i+1} , contributing a total of $2i-2n-2$. Meanwhile $w_4 w_3^{-1}$ has descents at x_i and x_{n-1} , and its inverse has descents at x_{n+1} and x_{2n-i} , contributing a total of $i+n-1-(n+1+2n-i) = 2i-2n-2$. If $x_{j+1} = 0$, w_b possesses another descent at x_n and w_b^{-1} does as well, cancelling out, whereas $w_4 w_3^{-1}$ possesses neither.

Since $\text{maj}(w_b) - \text{maj}(w_b^{-1})$ does not alter under this operation, and by proceeding down the two words performing such exchanges we will eventually cause them to match, any two words in the same CW_D make the same contribution to $\text{maj}(w) - \text{maj}(w^{-1})$. $\square$

Lemma 3 *If w_1 is in CW_D for $D = \{i_1, \dots, i_k\}$, in which words have length n and exactly j 1s, then the contribution of w_1 to $\text{maj}(w_1 w_2^{-1}) - \text{maj}(w_2 w_1^{-1})$ is $2j - 2(\text{maj}(\tau_1))$ for τ_1 any tableaux with descent set D .*

Proof Since all elements of any CW_D contribute the same amount to $\text{maj}(w) - \text{maj}(w^{-1})$, we can select one of the simpler ones and calculate what that contribution is. Let us examine the word in which 1s are entered as far to the right as possible:

$$\begin{array}{ccccccccc} [0 \cdots 0] & 01 & [0 \cdots 0] & 01 & \cdots & 01 & [1 \cdots 10 \cdots 0] & 01 & [1 \cdots 1] & 01 & \cdots & 01 & [1 \cdots 1] \\ l_1 & & l_2 & & & & l_g & & l_{g+1} & & & & l_{k+1} \end{array}$$

It is notationally convenient to use l_i to denote the lengths of the segments between 01 patterns rather than differences between the i_k counting backward from n . Note that any of these lengths may be 0, in which case l_g may be nonunique and any nearby empty segment may be assigned the label. If there is no segment in which 1s and 0s are mixed, the calculation that follows is similar.

What is the contribution this word makes to $\text{maj}(w_1 w_2^{-1}) - \text{maj}(w_2 w_1^{-1})$?

In $w_1 w_2^{-1}$, the first descent occurs at place $l_1 + 2$. The next occurs at place $l_1 + l_2 + 4$, and so forth up to $l_1 + \dots + l_{g-2} + 2(g-2)$. The descent following the segment labeled l_{g-1} is further ahead of this pattern by a distance equal to the number of 1s remaining after the k 1s in the 01 patterns are accounted for and the first segments of length l_{g+1} through l_{k+1} are filled: thus it occurs at $l_1 + \dots + l_{g-1} + 2(g-1) + (j-k - (l_{g+1} + \dots + l_{k+1}))$. The descent following the segment labeled l_g comes at the end of the following full segment of 1s, so it happens at place $l_1 + \dots + l_g + 2g + l_{g+1}$. This pattern holds up to the descent following the segment labeled l_{k-1} , which occurs after the full segment of 1s labeled l_k , thus at place $l_1 + \dots + l_{k-1} + 2(k-1) + l_k$. The segments labeled l_k and l_{k+1} do not have descents following them; as in the previous discussion, whether or not a descent occurs in the transition from the end of w_1 to the beginning of w_2^{-1} does not matter, since if one does it will be cancelled upon subtraction.

The sum of the places of descents in the w_1 half of w is thus

$$l_1(k-1) + l_2(k-2) + \dots + l_{k-1}(1) + (k-1)(k) + (j-k) - l_{k+1}.$$

In w^{-1} , the inverted form of w_1 appears after w_2 , thus:

$$(w_2) \underbrace{[0 \dots 0]}_{l_{k+1}} 01 \underbrace{[0 \dots 0]}_{l_k} 01 \dots 01 \underbrace{[1 \dots 10 \dots 0]}_{l_g} 01 \underbrace{[1 \dots 1]}_{l_{g-1}} 01 \dots 01 \underbrace{[1 \dots 1]}_{l_1}$$

and we note that w_2 is of length $n = l_1 + \dots + l_{k+1} + 2k$.

Here the first descent in this half of the word comes at place $n + l_{k+1} + 2$. The next occurs at $n + l_{k+1} + l_k + 4$, and so forth until we reach l_g . The number of 1s in this segment is the number of 0s in the segment's original state. Thus the relevant descent occurs at $n + l_{k+1} + \dots + l_{g+1} + 2(k-g+1) + l_g - (j-k - (l_{g+1} + \dots + l_{k+1}))$. The next descent occurs at place $n + l_{k+1} + \dots + l_g + 2(k-g+2) + l_{g-1}$, and so forth until the last descent occurs at place $n + l_{k+1} + \dots + l_3 + 2(k-1) + l_2$. Totaling, we find a contribution to $\text{maj}(w^{-1})$ of

$$\begin{aligned} & (n)(k-1) + (k-1)(k) + l_{k+1}(k) + \dots + l_2(1) - (j-k) \\ &= (l_1 + \dots + l_{k+1} + 2k)(k-1) + (k-1)(k) + l_{k+1}(k) + \dots + l_2(1) - (j-k). \end{aligned}$$

Subtract the latter contribution from the former: the total contribution of w_1 to $\text{maj}(w) - \text{maj}(w^{-1})$ is

$$\begin{aligned}
& 2(j - k) - 2k(k - 1) + l_1(0) + l_2(-2) + l_3(-4) + \cdots + l_{k+1}(-2k) \\
& = 2j - 2k^2 + l_1(0) + l_2(-2) + l_3(-4) + \cdots + l_{k+1}(-2k).
\end{aligned}$$

For the associated set of tableaux, it is straightforward to total up in terms of the l_i the maj statistic of a tableaux with descent set identified by this D , confirming it to be $l_{k+1}k + l_k(k - 1) + \cdots + l_2(1) + k^2$. Thus the contribution of w_1 to $\text{maj}(w) - \text{maj}(w^{-1})$ is $2j - 2(\text{maj}(\tau_1))$, as required.

□

Now we have that w_1 contributes $2j - 2(\text{maj}(\tau_1))$ to $\text{maj}(w) - \text{maj}(w^{-1})$, where $\text{maj}(\tau_1) = \text{maj}(\sigma^{-1})$. Likewise w_2 , in the opposite position, contributes $-(2j - 2(\text{maj}(\tau_2)))$, so that the total of the two contributions is $2(\text{maj}(\tau_2) - \text{maj}(\tau_1))$: exactly twice the difference between the maj statistics of each tableaux, which are the maj statistics of the corresponding permutations σ and σ^{-1} respectively. Summing over all Catalan half-words and corresponding tableaux, we have proved the theorem. □

4 Generalization

The next step would seem to be applying a similar map to 012 words and permutations in $S_n(1234)$ via tableaux of up to 3 columns. We would need to find the appropriate rewrite of Conjecture 1, determine the necessary rule for the generalized Catalan word, and work out the correct map. The larger goal, naturally, would be to prove Conjecture 1 fully by performing a similar procedure for the general case of $01 \dots k$ words and tableaux of up to $k + 1$ columns.

References

- [1] Chunwei Song, The limiting case of Haglund's q, t -Schröder theorem and an involution formula. *Discrete Mathematics* 308 (2008) 5218-5229.
- [2] Jim Haglund, A proof of the q, t -Schröder conjecture, *Internat. Math. Res. Not.* (11) (2004) 525-560.